To appear in ApJ Letters.
Preprint typeset using LaTeX style emulateapj v. 5/2/11

arXiv:1407.5344v2 [astro-ph.SR] 26 Jul 2014

SIMP J2154–1055: A NEW LOW-GRAVITY L4β BROWN DWARF CANDIDATE MEMBER OF THE ARGUS ASSOCIATION

Jonathan Gagné[1], David Lafrenière[1], René Doyon[1], Étienne Artigau[1], Lison Malo[1,2], Jasmin Robert[1] and Daniel Nadeau[1]

[1] Département de Physique, Université de Montréal, C.P. 6128 Succ. Centre-ville, Montréal, QC H3C 3J7, Canada and
[2] Canada-France-Hawaii Telescope, 65-1238 Mamalahoa Hwy, Kamuela, HI 96743, USA

To appear in ApJ Letters.

ABSTRACT

We present SIMP J21543454–1055308, a new L4β brown dwarf identified in the *Sondage Infrarouge de Mouvement Propre* survey that displays signs of low gravity in its near-infrared spectrum. Using the Bayesian Analysis for Nearby Young AssociatioNs II (BANYAN II), we show that it is a candidate member of the Argus association, albeit with a 21% probability that it is a contaminant from the field. Measurements of radial velocity and parallax will be needed to verify its membership. If it is a member of Argus (age 30–50 Myr), then this object would have a planetary mass of $10 \pm 0.5\ M_{\rm Jup}$.

Subject headings: brown dwarfs — methods: data analysis — proper motions — stars: kinematics and dynamics

1. INTRODUCTION

In the last decade, several brown dwarfs (BDs) in the field have been identified as displaying low-gravity features attributable to youth (Kirkpatrick et al. 2008; Cruz et al. 2009; Allers & Liu 2013; Liu et al. 2013; Gagné et al. 2014a). One could expect that a fraction of those young objects are unrecognized members of kinematic associations. However, the lack of parallax or radial velocity measurements for those objects prevents us from directly assessing their kinematics, which makes the identification of BD members to nearby young associations very hard. Identifying such BDs of known age would provide benchmarks to study the atmospheres of very low-mass BDs that are known to have features similar to those of the few directly imaged giant exoplanets known today (e.g., Chauvin et al. 2004; Marois et al. 2008; Lafrenière et al. 2008; Lagrange et al. 2009; Bowler et al. 2013; Rameau et al. 2013; Kuzuhara et al. 2013; Currie et al. 2014; Naud et al. 2014). In order to address this, Gagné et al. (2014b) developed the Bayesian Analysis for Nearby Young AssociatioNs II tool (BANYAN II), a statistical analysis based on BANYAN I (Malo et al. 2013) that uses a naive Bayesian classifier to identify low-mass star and BD candidates to membership in nearby young associations, even in the absence of radial velocity and parallax measurements.

SIMP J21543454–1055308 (SIMP J2154–1055 hereafter) is a new low-gravity BD that we discovered as part of the *Sondage Infrarouge de Mouvement Propre* (SIMP). We briefly describe the SIMP survey in Section 2, and then present a spectroscopic follow-up of SIMP J2154–1055 in Section 3. In Section 4, we show evidence that this BD displays signs of low gravity, by comparing its near-infrared (NIR) spectrum with low-gravity standards and by using the gravity classification scheme of Allers & Liu (2013). We finally use the BANYAN II tool to show that this young BD is a candidate Argus association member (Section 5).

2. THE SIMP SURVEY

The SIMP survey (Artigau et al. 2009) has been initiated in 2006 to identify new nearby BDs from their red optical-to-NIR colors and high proper motions, by obtaining a second J-band epoch for 30% of the sky (mostly in the Southern Hemisphere), 5–8 yr after the Two Micron All Sky Survey (2MASS; Skrutskie et al. 2006). This was achieved by using the CPAPIR camera (Artigau et al. 2004) on both the CTIO 1.5m telescope and the Observatoire du Mont-Mégantic 1.6m telescope. The SIMP survey typically reached an astrometric precision of $0\farcs15$ and a photometric depth of $J = 17$. Any source in the SIMP survey that satisfied at least one of three filters based on proper motion (μ) and $I-J$ color was selected for a spectroscopic follow-up: (1) $\mu > 100$ mas yr^{-1} and detected in I with $I-J > 3.5$; (2) $\mu > 100$ mas yr^{-1} and not detected in I such that $I-J > 3$; or (3) $\mu > 200$ mas yr^{-1} and not detected in I. I magnitudes were selected either from the *SuperCOSMOS Sky Survey* (Hambly et al. 2001), the Sloan Digital Sky Survey (York et al. 2000), or the Catalina Sky Survey (CSS; Larson et al. 1998), as available and in this order of preference. These criteria were designed to reject close-by M-type dwarfs as well as distant objects. A large number of new BD candidates have been identified this way, from which more than a hundred have been followed with NIR spectroscopy and confirmed as new M5–T3 very low-mass stars and BDs, most of them having spectral types later than L0. Among those, we have identified SIMP J2154–1055, an L4β BD displaying signs of low-gravity. A few other SIMP discoveries have been highlighted in Artigau et al. (2006) and Artigau et al. (2011), whereas remaining discoveries will be presented in an upcoming paper (J. Robert et al., in preparation).

3. SPECTROSCOPIC FOLLOW-UP

The NIR spectrum of SIMP J2154–1055 presented here was obtained on 2008 September 16 at the IRTF (program number 2008B054), using SpeX in the prism mode with the $0\farcs5$ slit ($R \sim 150$), covering the 0.8–2.5 μm range. The source was moved along the slit in an ABBA pattern, with a total of 10 exposures of 180 s to achieve a signal-to-noise ratio of $\sim$180 per resolution element. Raw exposures were reduced and combined using the SpeXtool package (Cushing et al. 2004), and telluric corrections were applied in a standard way (Vacca et al. 2003), using the A-type star HD 202990 observed im-

mediately before the target and at a similar airmass.

4. SIGNS OF LOW-GRAVITY

The NIR spectrum of SIMP J2154–1055 was visually compared with various NIR spectral templates to assign it a spectral type and identify peculiar features, following the method of K. Cruz et al. (in preparation; see also Cruz & Núñez 2007). Field templates were built for all spectral types in the M5–T8 range by median-combining high-quality spectra that showed no peculiarities nor signs of unresolved binarity in the DwarfArchives.[1] Low-gravity templates were built from known L0γ–L4γ BDs that were classified as low-gravity objects in the optical or the NIR (Cruz et al. 2009; Allers & Liu 2013; K. Cruz et al., in preparation). This comparison showed that SIMP J2154–1055 matches the L4 templates better than any other spectral types. Furthermore, the L4γ template is a better match than the field L4. The key spectral regions that differentiate between those two templates are; (1) the depth of the KI_J feature at $\sim$1.25 μm; (2) the level of the red end of the J band ($\sim$1.3 μm); (3) the triangular shape of the H band (1.5–1.8 μm); and (4) the slope of the K band plateau (2.15–2.3 μm). We show a comparison of SIMP J2154–1055 with the L4 and L4γ templates in Figure 1, with those four regions identified. The L4 and L4γ templates were built using five and two distinct spectra respectively.

Allers & Liu (2013) developed an NIR classification scheme to determine spectral types in a way that should not be sensitive to surface gravity, using visual classification supplemented by the H_2O indices, and then determined a gravity class using various spectroscopic indices sensitive to surface gravity. Objects for which most low-gravity indices are strong are classified as very low-gravity (VL-G) objects, and those having only a few indices as intermediate-gravity (INT-G) objects. Those without significant signs of youth are classified as field gravity. We have used this scheme to classify SIMP J2154–1055, with the exception that we used solely visual classification to determine its spectral type. Results are presented in Table 1, as well as in Figure 2. We find it is classified as an INT-G BD. Allers & Liu (2013) indicate that their INT-G and VL-G classifications respectively correspond to the β and γ classifications defined for optical spectra by Cruz et al. (2009). For this reason, we adopt L4β as the NIR spectral type of SIMP J2154–1055. Obtaining a higher-resolution NIR spectrum for this object would be useful in verifying that its alkali line equivalent widths are effectively weaker than normal. SIMP J2154–1055 is detected in the ALLWISE (Kirkpatrick et al. 2014) $W3$ channel, but does not display signs of infrared excess. We also show that it displays the reddest $J-K_s$ NIR color of all currently known L4 dwarfs (Figure 3), an effect likely attributed to thicker clouds in its photosphere. It is suspected that low gravity (youth) can cause such thicker clouds (Burgasser et al. 2008b).

5. ARGUS MEMBERSHIP

We used the BANYAN II tool (Gagné et al. 2014b) to verify if SIMP J2154-1055 is a candidate member of nearby young associations. We used the 2MASS and ALLWISE astrometry, as well as data from the DEep Near Infrared Survey of the Southern Sky (DENIS; Epchtein 1998) and measurements obtained with CPAPIR through the SIMP survey to calculate its proper motion (Table 1). We find that SIMP J2154–1055 has a 83.8% probability of being a member of the Argus association (30–50 Myr; Torres et al. 2008). The Argus association of stars was first identified by Makarov & Urban (2000), which proposed that the open cluster IC 2391 was a part of it. Torres et al. (2003) used the method of convergent point proper motion to show that both associations shared common kinematics, thus confirming this hypothesis. This association currently has 11 known A0–M5 bona fide members (Malo et al. 2013), from which the latest-type is the nearby star AP Col (Riedel et al. 2011). Gagné et al. (2014b) also proposed three low-gravity L-type BDs as candidate members to this association. Since the probability for the Argus membership of SIMP J2154-1055 is derived from a naive Bayesian classifier, it is expected to be biased when using dependent measurements as input observables (such as is the case here, see detail in Gagné et al. 2014b). Hence, we used the Monte Carlo contamination analysis presented in Gagné et al. (2014b) to obtain a probability of 20.5% that SIMP J2154–1055 is a young false-positive from the field. In the present case, the contamination and membership probabilities are almost exactly complementary, but this is not true in general.

In Figure 4, we show the proper motion of SIMP J2154–1055 compared with those of known bona fide members in Argus. Since members are spread across a large portion of the sky, the direction of their proper motion can be different, however it is expected that the great circles corresponding to the motion of all members of a given moving group will pass close to the group's apex and antapex. The proper motion of SIMP J2154–1055 thus seems consistent with a membership to Argus, since its great circle is closer to the apex than 6/11 bona fide members. Its very red $J-K_S$ colors (Figures 2 and 3) are also consistent with this interpretation. The BANYAN II tool allows us to predict that this object should have a distance of $22.1^{+2.8}_{-2.4}$ pc and a radial velocity of -13.0 ± 1.4 km s^{-1}

TABLE 1
PROPERTIES OF SIMP J2154–1055

Property	Value
R.A.	21:54:34.54
Decl.	–10:55:30.8
$\mu_\alpha \cos\delta$ (mas yr^{-1})[a]	169.2 ± 8.6
μ_δ (mas yr^{-1})[a]	-1.6 ± 8.8
I (CSS)[b]	21.63 ± 0.06
J (2MASS)	16.44 ± 0.12
H (2MASS)	15.07 ± 0.08
K_s (2MASS)	14.20 ± 0.07
$W1$ (ALLWISE)	13.36 ± 0.03
$W2$ (ALLWISE)	12.91 ± 0.03
$W3$ (ALLWISE)	12.54 ± 0.53
$W4$ (ALLWISE)	> 9.05
NIR spectral type	L4 $\beta \pm 0.5$
Gravity score[c]	*1021*
Gravity class[c]	INT-G
Estimated mass ($M_{\rm Jup}$)	10–11

[a] Proper motion from 2MASS, ALLWISE, SIMP and DENIS.
[b] Combined measurement from 34 epochs.
[c] See Allers & Liu (2013). Scores of *0*, *1* or *2* mean high, medium and low gravity, respectively. The four numbers are based on *FeH*, *VO*, *K I* and the *H*-band continuum shape, respectively.

[1] http://dwarfarchives.org



FIG. 1.— NIR spectrum of SIMP J2154-1055, compared with field (blue) and very low-gravity (red) template spectra of the same spectral type. Each band was normalized individually. We classify this object as L4β using the classification scheme of Allers & Liu (2013; Section 4), however we do not currently have access to any L4β BD to build a template and compare it with SIMP J2154–1055. The gray shaded region represents the scatter of the individual objects used to create the templates and the black vertical lines represent the measurement uncertainty on each bin of the observed spectrum. Beige shaded regions correspond to the locations of gravity-sensitive spectral indices defined by Allers & Liu (2013). The global continuum shape of this new BD is a better match to the L4γ template, despite its H-band continuum that seems to be an intermediate case between the two templates, which is consistent with our classification. We denote regions useful to differentiate between the field and low-gravity templates with purple numbers (Section 4).

if it is a member of Argus, or a distance of $23.7^{+6.4}_{-5.2}$ pc and a radial velocity of -8.9 ± 9.1 km s^{-1} if it is a field object. Using its 2MASS and ALLWISE apparent magnitudes, statistical distance, AMES-COND isochrones (Baraffe et al. 2003) and CIFIST2011 BT-SETTL atmosphere models (Allard et al. 2013) we determined that, at the age of Argus (30–50 Myr; Torres et al. 2008), SIMP J2154–1055 has a predicted mass of 10–11 $M_{\rm Jup}$. Measurements of radial velocity and parallax will be needed to assert its membership.

6. CONCLUSIONS

We present the discovery of a new young L4β BD, identified as part of the SIMP survey. We classify it as intermediate-gravity BD following the NIR gravity classification scheme of Allers & Liu (2013). We determine that it has an 84% probability of being an Argus member. We estimate that if it is a member of the Argus association with an adopted age of 30–50 Myr, its mass would be 10 ± 0.5 $M_{\rm Jup}$, in the planetary-mass regime. Adding this object to the currently known late-type low-mass stars and BDs will help shape our understanding of the properties of low-gravity, low-pressure atmospheres reminiscent of those of giant, gaseous exoplanets. The data presented in this Letter can be downloaded at our group's Web site.[2]

The authors would like to thank the anonymous referee for its suggestions, as well as Kelle Cruz and Jacqueline Faherty for useful comments and discussions and the sharing of spectroscopic templates. We thank Katelyn Allers for the sharing of several low-gravity spectra. This work was supported in part through grants from the Fonds de Recherche Québécois - Nature et Technologie and the Natural Science and Engineering Research Council of Canada. This research has benefited from the SpeX Prism Spectral Libraries, maintained by Adam Burgasser at `http://pono.ucsd.edu/~adam/browndwarfs/spexprism`, as well as the M, L, T and Y dwarf compendium housed at DwarfArchives.org. This research made use of; the SIMBAD database and VizieR catalogue access tool, operated at the Centre de Données astronomiques de Strasbourg, France; data products from the 2MASS, which is a joint project of the University of Massachusetts and the Infrared Processing and Analysis Center (IPAC)/California Institute of Technology (Caltech), funded by the National Aeronautics and Space Administration (NASA) and the National Science Foundation (Skrutskie et al. 2006); data products from the *Wide-field Infrared Survey Explorer*, which is a joint project of the University of California, Los Angeles, and the Jet Propulsion Laboratory (JPL)/Caltech, funded by NASA (Wright et al. 2010); the NASA/IPAC Infrared Science Archive, which is operated by JPL, Caltech, under contract with NASA; the Infrared Telescope Facility (IRTF), which is operated by the University of Hawaii under Cooperative Agreement NNX-08AE38A with NASA, Science Mission Directorate, Planetary Astronomy Program; and of tools provided by Astrometry.net. This publication uses observations obtained at IRTF through program number 2008B054. The authors recognize and acknowledge the very significant cultural role and reverence that the summit of Mauna Kea has always had within the indigenous Hawaiian community. We are most fortunate to have the opportunity to

[2] `http://www.astro.umontreal.ca/~gagne`



(a) FeH_Z (b) H_{CONT}

(c) KI_J (d) $J-K_S$

FIG. 2.— Panels (a)-(c): Gravity-sensitive spectroscopic indices for SIMP J2154–1055 (black star) compared with the field dwarf sequence (dark blue line) and its associated scatter (pink shaded region delimited by dashed pale blue lines). The dashed dark blue line delimits the intermediate-gravity and very low-gravity regimes. All objects classified as INT-G and VL-G in Allers & Liu (2013) are also displayed as green and purple dots, respectively.
Panel (d): 2MASS $J-K_S$ colors of SIMP J2154–1055 (black star) compared with the median values and standard deviation for field dwarfs (pink shaded region delimited by dashed pale blue lines; constructed from field BDs in the DwarfArchives), young field BDs (red dots; see Gagné et al. 2014b for the extensive list), and moderately young BD candidates (open purple symbols; Bannister & Jameson 2005; Jameson et al. 2008; Bihain et al. 2010; Casewell et al. 2014); circles for the Pleiades (∼125 Myr), downside triangles for Ursa Major (∼400 Myr), right triangles for Coma Berenices (∼500 Myr) and left triangles for the Hyades (∼625 Myr). The colors of SIMP J2154–1055 are consistent with its probable membership in Argus (Section 5).

conduct observations from this mountain.

REFERENCES

Allard, F., Homeier, D., Freytag, B., Schaffenberger, & Rajpurohit, A. S. 2013, MmSAI, 24, 128
Allers, K. N., & Liu, M. C. 2013, ApJ, 772, 79
Artigau, É., Doyon, R., Lafrenière, D., et al. 2006, ApJL, 651, L57
Artigau, É., Doyon, R., Vallée, P., Riopel, M., & Nadeau, D. 2004, Proc. SPIE, 5492, 1479
Artigau, É., Lafrenière, D., Doyon, R., et al. 2009, Cool Stars, 1094, 493
Artigau, É., Lafrenière, D., Doyon, R., et al. 2011, ApJ, 739, 48
Bannister, N. P., & Jameson, R. F. 2005, Astron. Nachr., 326, 1020
Baraffe, I., Chabrier, G., Barman, T. S., Allard, F., & Hauschildt, P. H. 2003, A&A, 402, 701
Bihain, G., Rebolo, R., Zapatero Osorio, M. R., Béjar, V. J. S., & Caballero, J. A. 2010, A&A, 519, 93
Bowler, B. P., Liu, M. C., Shkolnik, E. L., & Dupuy, T. J. 2013, ApJ, 774, 55
Burgasser, A. J., Cruz, K. K., Cushing, M. C., et al. 2010, ApJ, 710, 1142
Burgasser, A. J., Cruz, K. K., Liu, M. C., Ireland, M. J., & Dupuy, T. J. 2008a, ApJ, 681, 579
Burgasser, A. J., Looper, D. L., Kirkpatrick, D. J., Cruz, K. K., & Swift, B. J. 2008b, ApJ, 674, 451
Casewell, S. L., Littlefair, S. P., Burleigh, M. R., & Roy, M. 2014, MNRAS, 441, 2644
Chauvin, G., Lowrance, P., Lagrange, A.-M., et al. 2004, A&A, 425, L29
Cruz, K. K., Kirkpatrick, D. J., & Burgasser, A. J. 2009, AJ, 137, 3345
Cruz, K. K., Lowrance, P., Reid, N. I., Liebert, J., & Kirkpatrick, D. J. 2003, AJ, 126, 2421
Cruz, K. K., & Núñez, A. 2007, in Cool Stars 17, Poster Presentation



FIG. 3.— NIR spectrum of SIMP J2154–1055 (L4β) compared with other known L4 BDs (Reid et al. 2000; Cruz et al. 2003; Kendall et al. 2004; Cruz et al. 2007; Reid et al. 2008; Burgasser et al. 2008a; Lafrenière et al. 2008; Lafrenière et al. 2010; Burgasser et al. 2010; Allers & Liu 2013). Although it is classified as INT-G, it is the reddest L4 BD yet identified. All spectra are normalized to their median in the 1.27–1.33 μm range.

Cruz, K. K., Reid, N. I., Kirkpatrick, D. J., et al. 2007, AJ, 133, 439
Currie, T., Daemgen, S., Debes, J., et al. 2014, ApJL, 780, L30
Cushing, M. C., Vacca, W. D., & Rayner, J. T. 2004, PASP, 116, 362
Epchtein, N. 1998, New Horizons from Multi-Wavelength Sky Surveys, 179, 106
Gagné, J., Faherty, J. K., Cruz, K. K., et al. 2014a, ApJL, 785, L14
Gagné, J., Lafrenière, D., Doyon, R., Malo, L., & Artigau, É. 2014, ApJ, 783, 121
Hambly, N. C., MacGillivray, H. T., Read, M. A., et al. 2001, MNRAS, 326, 1279
Jameson, R. F., Lodieu, N., Casewell, S. L., Bannister, N. P., & Dobbie, P. D. 2008, MNRAS, 385, 1771
Kendall, T. R., Delfosse, X., Martín, E. L., & Forveille, T. 2004, A&A, 416, L17
Kirkpatrick, D. J., Lowrance, P., Cruz, K. K., et al. 2008, ApJ, 689, 1295
Kirkpatrick, D. J., Schneider, A. C., Fajardo-Acosta, S., et al. 2014, ApJ, 783, 122
Kuzuhara, M., Tamura, M., Kudo, T., et al. 2013, ApJ, 774, 11
Lafrenière, D., Jayawardhana, R., & van Kerkwijk, M. H. 2008, ApJL, 689, L153
Lafrenière, D., Jayawardhana, R., & van Kerkwijk, M. H. 2010, ApJ, 719, 497
Lagrange, A.-M., Gratadour, D., Chauvin, G., et al. 2009, A&A, 493, L21
Larson, S., Brownlee, J., Hergenrother, C., & Spahr, T. 1998, BAAS, 30, 1037
Lawrence, A., Warren, S. J., Almaini, O., et al. 2007, MNRAS, 379, 1599
Liu, M. C., Magnier, E. A., Deacon, N. R., et al. 2013, ApJL, 777, L20
Makarov, V. V., & Urban, S. 2000, MNRAS, 317, 289
Malo, L., Doyon, R., Lafrenière, D., et al. 2013, ApJ, 762, 88
Marois, C., Macintosh, B., Barman, T. S., et al. 2008, Sci, 322, 1348
Naud, M.-E., Artigau, É., Malo, L., et al. 2014, ApJ, 787, 5
Rameau, J., Chauvin, G., Lagrange, A.-M., et al. 2013, ApJL, 779, L26
Reid, N. I., Kirkpatrick, D. J., Gizis, J. E., et al. 2000, AJ, 119, 369
Reid, N. I., Lowrance, P., Cruz, K. K., et al. 2008, AJ, 136, 1290
Riedel, A. R., Murphy, S. J., Henry, T. J., et al. 2011, AJ, 142, 104
Skrutskie, M. F., Gizis, J. E., Kirkpatrick, D. J., et al. 2006, AJ, 131, 1163
Torres, C. A. O., Quast, G. R., de La Reza, et al. 2003, in Open Issues in Local Star Formation, ed. J. Lépine & J. Gregorio-Hetem (Astrophysics and Space Science Library, Vol. 299; Ouro Preto, Brazil; Dordrecht: Kluwer), 83
Torres, C. A. O., Quast, G. R., Melo, C. H. F., & Sterzik, M. F. 2008, Young Nearby Loose Associations (The Southern Sky ASP Monograph Publications; San Francisco, CA, Vol. I: ASP)
Vacca, W. D., Cushing, M. C., & Rayner, J. T. 2003, PASP, 115, 389
Wright, E. L., Eisenhardt, P. R. M., Mainzer, A. K., et al. 2010, AJ, 140, 1868
York, D. G., Adelman, J., Anderson, J. E. J., et al. 2000, AJ, 120, 1579



FIG. 4.— Position and proper motion (red arrow) of SIMP J2154–1055 with its associated projected great circle on the celestial sphere (pink line), as compared with bona fide members in the Argus association (green lines and arrows). It can be seen that the proper motion of SIMP J2154–1055 is consistent with other bona fide members: its projected great circle falls closer to Argus' apex and antapex (blue circles) than 6 out of 11 known members. The solar motion's apex and antapex are displayed with a black plus sign and cross, respectively.